

NAUMAN QUAMAR, BDS, MS
— Specialist in Periodontics & Implantology —

_____ (ofc) 469.231.1155 (cell) drq@vipsperio.com www.vipsperio.com

POST-OPERATIVE INSTRUCTIONS FOLLOWING DENTAL SURGERY

THINGS TO EXPECT:

- Bleeding:** Bleeding or "oozing" for the first 12 to 24 hours.
Swelling: This is normal following a surgical procedure in the mouth. It should reach its maximum in two-to-three days and should begin to diminish by the fifth post-operative day.
Discomfort: The most discomfort that you may experience may occur for a few hours after the sensation returns to your mouth. It may gradually increase again for 2-3 days, then begin to diminish over the next few days.

THINGS TO DO IMMEDIATELY FOLLOWING SURGERY:

- Bleeding:** Place gauze over surgical sites and maintain pressure by biting for at least one hour. Repeat as needed. Keep head elevated, and rest. Do not suck or spit excessively. (Also, please refrain from blowing into musical instruments.)
NOTE: Some "oozing" and discoloration of saliva is normal. If bleeding persists, replace the gauze with a clean folded gauze placed over the surgical site, and maintain the pressure until the bleeding stops.
- Swelling:** Place ice or cold compresses on the region of surgery for ten minutes every half-hour for the first eight to 12 hours.
NOTE: Ice bags or cold compresses should be used only on the day of surgery.
- Smoking:** Avoid smoking during the healing period.
- Discomfort:** Take medications as directed for **PAIN**. Mild-to-moderate pain can be relieved by non-prescription Advil, Aleve, or Orudis. For more severe pain, take the prescription pain medication as directed. Remember that these medications can take up to 30 minutes to one hour to take effect. If you are using any of these medications for the first time, exercise caution with the initial doses (start with ½ a pill).
- Diet:** A nutritious liquid or soft diet will be necessary for the first weeks after surgery. Healing will occur in weekly increments; therefore, it is best to **gradually** (in weekly increments) return the diet and/or other mouth/oral activities back to normal.
- Physical Activity:** For the first 24 to 48 hours, one should **REST**. Patients who have sedation should refrain from driving an automobile or from engaging in any task that requires alertness for the next 24 hours.

THE DAYS AFTER SURGERY:

1. Brush teeth carefully.
2. Beginning 24 hours after the surgery, rinse mouth with **WARM SALT WATER** (or prescription mouth rinse). Continue rinsing three-to-five times per day for seven days, then begin irrigating per dentist's instructions (see #7 below).
3. If **ANTIBIOTICS** are prescribed, be **SURE** to take **ALL** that have been prescribed, **AS DIRECTED**.
4. Use **WARM, MOIST HEAT** on face for swelling, if any. Continue until the swelling subsides. A warm, wet washcloth or heating pad will suffice.
5. If **SUTURES** were used, they will dissolve on their own.
6. **RETURN TO YOUR DENTIST'S OFFICE** five-to-seven days after the surgery for irrigation instructions.

CONTACT THE DOCTOR IF:

1. Bleeding is excessive and cannot be controlled.
2. Discomfort is poorly controlled.
3. Swelling is excessive, spreading, or continuing to enlarge after 60 hours.
4. Allergic reactions to medications occur, which are causing a generalized rash or excessive itching.

CONTACT EMERGENCY MEDICAL SERVICES ("EMS") OR CALL "911" IF:

Patient loses or has lost consciousness.

**** BE SURE TO CHECK THE WEBSITE FOR ADDITIONAL INFORMATION ****

— ***www.vipsperio.com*** —